



Clinical Pharmacist Practitioner Freelance Writer US Pharm. 2023;48 (6):17-21. ABSTRACT: Testosterone is the treatment of choice for testosterone deficiency (defined as low serum testosterone combined with symptoms of low testosterone) and hypogonadism (low testosterone with infertility).



 **BUY STEROIDS ONLINE** 

Switching between testosterone enanthate (Delatestryl) and testosterone .



Pros • Works well to replace testosterone in your body • Applied once a day to your skin, so it's easy to use • Available in tubes, packets, or as a pump • Some brands are available as lower-cost generics Cons • Can cause skin irritation • Need to avoid skin-to-skin transfer to other people • Need to wait hours before you can shower, swim, or bathe

Delatestryl (Testosterone Enanthate): Side Effects, Uses . - RxList



Depo-Testosterone (Pfizer Inc, New York, NY, USA) is the most cost-efficient TRT option and is only administered every 2 weeks, which may improve patient compliance, although the dose interval can be adjusted to once per week. Androderm is the most efficacious TRT medication, with up to 92% of patients reaching normal testosterone levels.

Testosterone Therapy With Subcutaneous Injections: A Safe, Practical .



DELATESTRYL (testosterone enanthate) should not be used to treat non-specific symptoms suggestive of hypogonadism if testosterone deficiency has not been demonstrated and if other etiologies responsible for the symptoms have not been excluded.

Depo-Testosterone vs Testosterone Comparison - Drugs



Injectable Testosterone . Depo-testosterone (testosterone cypionate). Depo-testosterone is the most popularly prescribed TRT medication in the US. 17. and is a long-acting testosterone commonly prescribed once every 2-weeks. The most common dose is 200 milligrams (mg) in 1 milliliter (ml) 18. although it is available in two doses: 100 mg/ml and .

Pharmacology of testosterone replacement therapy preparations


TABLE 2
FDA-approved testosterone formulations²

Formulation	Trade names	Dosing, route, and administration	Benefits	Potential adverse effects or drawbacks
Testosterone cypionate, testosterone enanthate		75-100 mg IM every week or 150-200 mg IM every 2 weeks	Short-acting preparation that allows for drug withdrawal in the event of adverse effects	Fluctuation of testosterone levels
Testosterone undecanoate	Aveed	750 mg IM followed in 4 weeks by another 750 mg IM, then 750 mg every 10 weeks thereafter	Steady-state testosterone levels without fluctuation	Long-acting formulation; does not allow for rapid discontinuation of therapy if adverse effects develop
Transdermal and transmucosal testosterone	Androderm (patch)	A single, 2-6-mg patch daily	Most closely mirrors physiologic testosterone levels	May cause skin irritation; risk of transfer to partner or children
	Axiron (axillary)	30-120 mg daily		
	AndroGel, Testim, Vogelxo (gel)	50-100 mg daily applied to shoulders, arms, or abdomen		
	Fortesta (gel)	10-70 mg daily applied to thighs		
	Natesto (intranasal)	11 mg 3 times daily		
	Striant (buccal)	30 mg every 12 hours		
Oral testosterone	Androxy (fluoxymesterone)	5-20 mg daily in a single or divided doses	Daily oral dosing	Difficult to obtain normal physiologic levels; risk of hepatotoxicity
	Android, Methitest, Testred (methyltestosterone)	10-50 mg daily		
Subdermal testosterone pellets	Testopel	150-450 mg implanted subcutaneously every 3-6 months	Long duration of action	Infection at implantation site; explantation

IM, intramuscular.

Testosterone cypionate (Depo-Testosterone) is an injectable form of testosterone that's used to treat low testosterone in adult males. It's injected into your buttock muscle and lasts for a long time, so it only needs to be used once or twice a month. But it might raise the risk of serious side effects including heart attack and stroke .

Testosterone (Intramuscular Route, Subcutaneous Route)



Depo- Testosterone (testosterone cypionate injection) is an androgen indicated for replacement therapy in the male in conditions associated with symptoms of deficiency or absence of endogenous testosterone including, 1) Primary hypogonadism (congenital or acquired) - testicular failure due to cryptorchidism, bilateral torsion, orchitis, vanishi.

Depo-Testosterone (Testosterone Cypionate Injection): Side . - RxList



Descriptions Testosterone injection is used for the treatment of men whose bodies do not make enough natural testosterone, a condition called hypogonadism. Testosterone is a male hormone responsible for the growth and development of the male sex organs and maintenance of secondary sex characteristics.







Compare AndroGel vs. Depo-Testosterone - GoodRx

Compare
AndroForte®
 to **AndroGel®**



VS




	AndroForte®	AndroGel®
Strength 	5%	1.62%
Doses per Pack 	50	30
Number of Days 	50	30
Initial Dose 	1 ml of cream delivers 50mg Testosterone	2 pumps of gel delivers 40.5mg Testosterone
Cost per Day 	\$3.30	\$14.83
Cost as advertised on  goodrx.com	\$165.00	\$445.00

*AndroGel price taken from GoodRX January 2015

• 1 yr. ago Timely-Bed656 Delatestryl vs. Depo-testosterone? Discussion Hello! I have been on T for 4 months now and I have been using/prescribed Delatestryl. However, my other ftm friends have been using depo-testosterone. Just wondering what the difference is between the two if there are any?? Physically or any faster changes etc?? TIA:)

PDF Testosterone injections are used in clinical practice as intramuscular .



TC 35 (Depo® -Testosterone) 34 200 mg/2 weeks: . 33 The three IM formulations that are approved by the FDA for use as TTh are testosterone cypionate (TC), TE, and testosterone undecanoate (TU). Depending on the formulation and dosage, following IM injection of TE or TC, supraphysiologic levels occur within a week after administration .

Comparison of Testosterone Replacement Therapy Medications for .


TYPES OF TESTOSTERONE MEDICATIONS

Trade Name	Generic	Route	Available Dosing	Common Dose	Common Frequency	Average Cost/Month (calculated from common dose)
Injectable						
depo-testosterone	testosterone cypionate	IM	100 mg/ml 200 mg/ml	200 mg/ml ¹³	Q 2 weeks	100mg/ml (10ml vial) = \$22.70 ¹³ 200mg/ml (10ml vial) = \$18.45 ¹³ 200mg/ml (1ml vial) = \$98.07 ¹³
Delatestryl	testosterone enanthate	IM	200 mg/ml	200 mg/ml ¹⁷	Q 2 weeks	200mg/ml (5ml vial) = \$23.46 ¹⁷
Nasal						
Natesto	N/A	Nasal gel	5.5 mg each actuation	11 mg (1 actuation per nostril) ¹¹	TID	3 bottles (30 days) = \$699.21 ¹¹
Buccal						
Striant	N/A	Buccal	30 mg	30 mg ¹²	BID	60 tablets = \$724.77 ¹²
Transdermal Patch						
Androderm	N/A	Patch	2 mg 4 mg	4 mg ²²	QD	30 patches = \$195.03 ¹⁹
Transdermal Gel Pump						
AndroGel 1% or 1.62%	testosterone 1% testosterone 1.62%	Gel pump	1%-12.5mg/1.25g 1.62%-20.25mg/1.25g	50mg (4 pumps) ¹⁹ 40.5mg (2 pumps) ¹⁹	QD	2 bottles 1% (30 days) \$309.24 ¹⁷ 1 bottle 1.62% (30 days) \$510.77 ¹⁷
Axiiron 2%	N/A	Gel pump	30mg /1.5ml	60 mg (2 pumps) ¹⁹	QD	1 bottle (30 days) \$630.78 ¹⁹
Fortesta 2%	testosterone 2%	Gel pump	10mg/0.5g	40mg (4 pumps) ²	QD	1 bottle (30 days) = \$281.80 ¹⁷
Vogelxo 1%	testosterone 1%	Gel pump	12.5 mg/1.25g	50mg (4 pumps) ¹⁷	QD	2 bottles (30 days) = \$283.69 ¹⁷
Transdermal Gel Packets						
AndroGel 1% or 1.62%	testosterone 1% testosterone 1.62%	Gel packet	1%-25mg/2.5g 1%-50mg/5g	50mg (1 packet) ¹⁷	QD	30 packets 1% = \$300.25 ¹⁷
Vogelxo 1%	testosterone 1%	Gel packet	1.62%-40.5mg/2.5g 1%-25mg/2.5g 1%-50mg/5g	40.5mg (1 packet) ¹⁹ 50mg (1 packet) ¹⁹	QD	30 packets 1.62% = \$516.41 ¹⁷ 30 packets 1% = \$270.30 ¹⁷


Medical conditions which necessitate testosterone replacement therapy (TRT) include primary hypogonadism, in which the condition originates in the testes, and secondary/hypogonadotropic hypogonadism, a disease of the hypothalamus or pituitary gland.



Compare Methyltestosterone vs. Depo-Testosterone - GoodRx

**TESTOSTERONE LEVELS:
NORMAL VS OPTIMAL**



TESTOSTERONE



Published on February 10, 2023 Key takeaways: Testosterone injections are effective medications for combating low testosterone ("T") levels. They're a common type of testosterone replacement therapy.

PDF Prescribing Information

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use OZOBAX® safely and effectively. See full prescribing information for OZOBAX.

OZOBAX (baclofen) oral solution
Initial U.S. Approval: 1977

INDICATIONS AND USAGE

- OZOBAX is a gamma-aminobutyric acid (GABA-ergic) agonist indicated for the treatment of spasticity resulting from multiple sclerosis, particularly for the relief of flexor spasms and concomitant pain, clonus, and muscular rigidity. (1)
- OZOBAX may also be of some value in patients with spinal cord injuries and other spinal cord diseases. (1)

Limitations of Use

OZOBAX is not indicated in the treatment of skeletal muscle spasm resulting from rheumatic disorders (1)

DOSAGE AND ADMINISTRATION

- Initiate OZOBAX with a low dosage, preferably in divided doses, administered orally. Increase gradually based on clinical response and tolerability. (2.1)
- The maximum dosage is 80 mg daily (20 mg four times a day). (2.1)
- When discontinuing, reduce the dosage slowly. (2.2)

DOSAGE FORMS AND STRENGTHS

Oral Solution: 5 mg/5 mL (3)

CONTRAINDICATIONS

- Hypersensitivity to baclofen (4)

WARNINGS AND PRECAUTIONS

- Abrupt discontinuation of baclofen has resulted in serious adverse reactions including death; therefore, reduce the dosage slowly when OZOBAX is discontinued. (5.1)
- Neonatal withdrawal symptoms can occur; gradually reduce the dosage and discontinue OZOBAX before delivery. (5.2)
- OZOBAX can cause drowsiness and sedation. Patients should avoid the operation of automobiles or other dangerous machinery until they know how the drug affects them. Advise patients that the central nervous system effects of OZOBAX may be additive to those of alcohol and other CNS depressants. (5.3)
- OZOBAX can cause exacerbation of the following: psychotic disorders, schizophrenia, or confusional states; autonomic dysreflexia; epilepsy. Use with caution in patients with these conditions (5.5, 5.6, 5.7)

ADVERSE REACTIONS

- The most common (up to 15% or more) adverse reactions in patients were drowsiness, dizziness, and weakness. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Melacel Pharmaceuticals, LLC at 1-833-469-6229 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

USE IN SPECIFIC POPULATIONS

- Pregnancy: Based on animal data, may cause fetal harm (8.1)
- Because baclofen is excreted unchanged through the kidneys it may be necessary to reduce the dosage in patients with impaired renal function. (8.6)

See 17 for PATIENT COUNSELING INFORMATION

Revised: 05/2020

FULL PRESCRIBING INFORMATION: CONTENTS*

1	INDICATIONS AND USAGE
2	DOSAGE AND ADMINISTRATION
2.1	Recommended Dosage
2.2	Discontinuation of OZOBAX
3	DOSAGE FORMS AND STRENGTHS
4	CONTRAINDICATIONS
5	WARNINGS AND PRECAUTIONS
5.1	Adverse Reactions From Abrupt Withdrawal of OZOBAX
5.2	Neonatal Withdrawal Symptoms
5.3	Drowsiness and Sedation
5.4	Poor Tolerability in Stroke Patients
5.5	Exacerbation of Psychotic Disorders, Schizophrenia, or Confusional States
5.6	Exacerbation of Autonomic Dysreflexia
5.7	Exacerbation of Epilepsy
5.8	Posture and Balance Effects
5.9	Ovarian Cysts
6	ADVERSE REACTIONS
6.1	Clinical Trials Experience
7	DRUG INTERACTIONS
7.1	CNS Depressants and Alcohol

8 USE IN SPECIFIC POPULATIONS

8.1	Pregnancy
8.2	Lactation
8.4	Pediatric Use
8.5	Geriatric Use
8.6	Renal Impairment
10	OVERDOSAGE
10.1	Symptoms of Baclofen Overdose
10.2	Treatment for Overdose
11	DESCRIPTION
12	CLINICAL PHARMACOLOGY
12.1	Mechanism of Action
12.2	Pharmacodynamics
12.3	Pharmacokinetics
13	NONCLINICAL TOXICOLOGY
13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility
14	CLINICAL STUDIES
16	HOW SUPPLIED/STORAGE AND HANDLING
16.1	How Supplied
16.2	Storage and Handling
17	PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

OZOBAX is indicated for the treatment of spasticity resulting from multiple sclerosis, particularly for the relief of flexor spasms and concomitant pain, clonus, and muscular rigidity.

OZOBAX may also be of some value in patients with spinal cord injuries and other spinal cord diseases.

Limitations of Use

OZOBAX is not indicated in the treatment of skeletal muscle spasm resulting from rheumatic disorders.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

Initiate OZOBAX with a low dosage, preferably in divided doses, administered orally. The following gradually increasing dosage regimen is suggested, but should be adjusted based on clinical response and tolerability:

5 mL (5 mg) three times a day for three days

10 mL (10 mg) three times a day for three days

15 mL (15 mg) three times a day for three days

20 mL (20 mg) three times a day for three days

Additional increases may be necessary up to the maximum recommended dosage of 80 mg daily (20 mg four times a day).

2.2 Discontinuation of OZOBAX

When discontinuing OZOBAX, reduce the dosage slowly and avoid abrupt withdrawal from the drug to help minimize the risk of adverse reactions (see Warnings and Precautions § 5.1).

3 DOSAGE FORMS AND STRENGTHS

Oral Solution: 5 mg/5 mL baclofen as a clear, colorless solution with a grape aroma.

4 CONTRAINDICATIONS

OZOBAX is contraindicated in patients with hypersensitivity to baclofen.

5 WARNINGS AND PRECAUTIONS

5.1 Adverse Reactions from Abrupt Withdrawal of OZOBAX

Abrupt discontinuation of baclofen, regardless of the cause, has resulted in adverse reactions that include hallucinations, seizures, high fever, altered mental status, exaggerated rebound spasticity, and muscle rigidity, that in rare cases has advanced to rhabdomyolysis, multiple organ-system failure, and death. Therefore, reduce the dosage slowly when OZOBAX is discontinued, unless the clinical situation justifies a rapid withdrawal.

5.2 Neonatal Withdrawal Symptoms

Withdrawal symptoms in neonates whose mothers were treated with oral baclofen throughout pregnancy have been reported starting hours to days after delivery. The symptoms of withdrawal in these infants have included increased muscle tone, tremor, jitteriness, and seizures. If the potential benefit justifies the potential risk to the fetus and OZOBAX is continued during pregnancy, gradually reduce the dosage and discontinue OZOBAX before delivery. If slow withdrawal is not feasible, advise the parents or caregivers of the exposed neonate of the potential for neonatal withdrawal.

5.3 Drowsiness and Sedation

Drowsiness and sedation have been reported in up to 60% of patients taking baclofen; the active ingredient in OZOBAX (see Adverse Reactions § 6.1). Patients should avoid operation of automobiles or other dangerous machinery and activities made hazardous by decreased alertness when starting OZOBAX or increasing the dose until they know how the drug affects them. Advise patients that the central nervous system depressant effects of OZOBAX may be additive to those of alcohol and other CNS depressants.

5.4 Poor Tolerability in Stroke Patients

OZOBAX should be used with caution in patients who have had a stroke. Baclofen has not significantly benefited patients with stroke. These patients have also shown poor tolerability to the drug.

5.5 Exacerbation of Psychotic Disorders, Schizophrenia, or Confusional States

OZOBAX should be used with caution in patients suffering from psychotic disorders, schizophrenia, or confusional states. If treated with OZOBAX, these patients should be kept under careful surveillance because exacerbations of these conditions have been observed with oral baclofen administration.


5.6 Exacerbation of Autonomic Dysreflexia

OZOBAX should be used with caution in patients with a history of autonomic dysreflexia. The presence of noxious stimuli or abrupt withdrawal of OZOBAX may cause an autonomic dysreflexic episode.

Delatestryl (testosterone enanthate) is an androgen, a male sex hormone, used in men who do not make enough of a natural substance called testosterone. Delatestryl is available in generic form. What Are Side Effects of Delatestryl? Common side effects of Delatestryl include: nausea, vomiting, headache,

anxiety, depression, skin color changes,

Pharmacokinetics of testosterone therapies in relation to diurnal .



info Specialty Drug testosterone cypionate Depo-Testosterone Used for Testosterone Replacement Testosterone cypionate (Depo-Testosterone) is an injectable form of testosterone that's used to treat low testosterone in adult males. It's injected into your buttock muscle and lasts for a long time, so it only needs to be used once or twice a month.

Antares' Xyosted Is One Of The Best Testosterone . - Seeking Alpha



Depo-Testosterone (Pfizer Inc, New York, NY, USA) is the most cost-efficient TRT option and is only administered every 2 weeks, which may improve patient compliance, although the dose interval can be adjusted to once per week. Androderm is the most efficacious TRT medication, with up to 92% of patients reaching normal testosterone levels.

Comparison of Testosterone Replacement Therapy Medications in the .

TYPES OF TESTOSTERONE MEDICATIONS						
Trade Name	Generic	Route	Available Dosing	Common Dose	Common Frequency	Average Cost/Month (calculated from common dose)
Injectable						
depo-testosterone	testosterone cypionate	IM	100 mg/ml 200 mg/ml	200 mg/ml ¹³	Q 2 weeks	100mg/ml (10ml vial) = \$22.70 ¹³ 200mg/ml (10ml vial) = \$18.45 ¹³ 200mg/ml (1ml vial) = \$98.07 ¹³
Delatestryl	testosterone enanthate	IM	200 mg/ml	200 mg/ml ¹⁷	Q 2 weeks	200mg/ml (5ml vial) = \$23.46 ¹⁷
Nasal						
Natesto	N/A	Nasal gel	5.5 mg each actuation	11 mg (1 actuation per nostril) ¹³	TID	3 bottles (30 days) = \$699.21 ¹⁷
Buccal						
Striant	N/A	Buccal	30 mg	30 mg ¹³	BID	60 tablets = \$724.77 ¹³
Transdermal Patch						
Androderm	N/A	Patch	2 mg 4 mg	4 mg ²²	QD	30 patches = \$195.03 ¹³
Transdermal Gel Pump						
AndroGel 1% or 1.62%	testosterone 1% testosterone 1.62%	Gel pump	1%-12.5mg/1.25g 1.62%-20.25mg/1.25g	50mg (4 pumps) ¹⁹ 40.5mg (2 pumps) ¹⁹	QD	2 bottles 1% (30 days) \$309.24 ¹⁷ 1 bottle 1.62% (30 days) \$510.77 ¹⁷
Axiron 2%	N/A	Gel pump	30mg /1.5ml	60 mg (2 pumps) ¹⁹	QD	1 bottle (30 days) \$630.78 ¹⁹
Fortesta 2%	testosterone 2%	Gel pump	10mg/0.5g	40mg (4 pumps) ⁷	QD	1 bottle (30 days) = \$281.80 ¹⁷
Vogelxo 1%	testosterone 1%	Gel pump	12.5 mg/1.25g	50mg (4 pumps) ¹⁷	QD	2 bottles (30 days) = \$283.69 ¹⁷
Transdermal Gel Packets						
AndroGel 1% or 1.62%	testosterone 1% testosterone 1.62%	Gel packet	1%-25mg/2.5g 1%-50mg/5g	50mg (1 packet) ¹⁹	QD	30 packets 1% = \$309.25 ¹⁷
1.62%	testosterone 1.62%	Gel packet	1.62%-20.25mg/1.25g 1.62%-40.5mg/2.5g	40.5mg (1 packet) ¹⁹		30 packets 1.62% = \$516.41 ¹⁷
Vogelxo 1%	testosterone 1%	Gel packet	1%-25mg/2.5g 1%-50mg/5g	50mg (1 packet) ¹⁹	QD	30 packets 1% = \$270.30 ¹⁷

Switching between testosterone enanthate (Delatestryl) and testosterone cypionate (Depo T) : r/ftm r/ftm r/ftm • 2 yr. ago [deleted] Switching between testosterone enanthate (Delatestryl) and testosterone

cypionate (Depo T) Discussion For those of you have switched from one to the other, what differences (if any) did you notice?

Comparison of Testosterone Replacement Therapy Medications in the .

TYPES OF TESTOSTERONE MEDICATIONS

Trade Name	Generic	Route	Available Dosing	Common Dose	Common Frequency	Average Cost/Month (calculated from common dose)
Injectable						
depo-testosterone	testosterone cypionate	IM	100 mg/ml 200 mg/ml	200 mg/ml ¹⁹	Q 2 weeks	100mg/ml (10ml vial) = \$22.70 ¹⁹ 200mg/ml (10ml vial) = \$18.45 ¹⁹ 200mg/ml (1ml vial) = \$98.07 ¹⁹
Delatestryl	testosterone enanthate	IM	200 mg/ml	200 mg/ml ¹⁷	Q 2 weeks	200mg/ml (5ml vial) = \$23.46 ¹⁷
Nasal						
Natesto	N/A	Nasal gel	5.5 mg each actuation	11 mg (1 actuation per nostril) ¹³	TID	3 bottles (30 days) = \$699.21 ¹³
Buccal						
Striant	N/A	Buccal	30 mg	30 mg ¹⁸	BID	60 tablets = \$724.77 ¹⁸
Transdermal Patch						
Androderm	N/A	Patch	2 mg 4 mg	4 mg ²²	QD	30 patches = \$195.03 ¹⁹
Transdermal Gel Pump						
AndroGel 1% or 1.62%	testosterone 1% testosterone 1.62%	Gel pump	1%-12.5mg/1.25g 1.62%-20.25mg/1.25g	50mg (4 pumps) ¹⁹ 40.5mg (2 pumps) ¹⁹	QD	2 bottles 1% (30 days) \$309.24 ¹³ 1 bottle 1.62% (30 days) \$510.77 ¹⁷
Axiron 2%	N/A	Gel pump	30mg /1.5ml	60 mg (2 pumps) ¹⁹	QD	1 bottle (30 days) \$630.78 ¹⁹
Fortesta 2%	testosterone 2%	Gel pump	10mg/0.5g	40mg (4 pumps) ²	QD	1 bottle (30 days) = \$281.80 ¹⁷
Vogelxo 1%	testosterone 1%	Gel pump	12.5 mg/1.25g	50mg (4 pumps) ¹⁷	QD	2 bottles (30 days) = \$283.69 ¹⁷
Transdermal Gel Packets						
AndroGel 1% or 1.62%	testosterone 1% testosterone 1.62%	Gel packet	1%-25mg/2.5g 1%-50mg/5g	50mg (1 packet) ¹⁷	QD	30 packets 1% = \$309.25 ¹⁷
1.62%	testosterone 1.62%	Gel packet	1.62%-20.25mg/1.25g 1.62%-40.5mg/2.5g	40.5mg (1 packet) ¹⁹		30 packets 1.62% = \$516.41 ¹⁷
Vogelxo 1%	testosterone 1%	Gel packet	1%-25mg/2.5g 1%-50mg/5g	50mg (1 packet) ¹⁹	QD	30 packets 1% = \$270.30 ¹⁷


LIBIDO AND ERECTILE FUNCTION A common indication for testosterone therapy is treatment of decreased sexual desire or erectile dysfunction. A systematic review found 23 randomized trials of.

Testosterone cypionate (Depo-Testosterone): Uses, Side Effects, Dosage .



A) Depo-Testosterone (testosterone cypionate intramuscular injection, generic): Up to 400 mg administered subcutaneously or intramuscularly every 1 to 4 weeks, not to exceed 400 mg every 2 weeks; OR B) Delatestryl (testosterone enanthate intramuscular injection, generic): Up to 400 mg administered subcutaneously or intramuscularly every 1 to 4 weeks, not to exceed 400 mg every 2 weeks; OR

A Review of Testosterone Therapy Options for Men - U. S. Pharmacist



testosterone cypionate intramuscular. TESTOSTERONE - INTRAMUSCULAR (tess-TOSS-ter-own)
COMMON BRAND NAME(S): Delatestryl, Tesamone. USES: This medication is used in men who do not make enough of a natural substance called testosterone. In males, testosterone is responsible for many normal functions, including growth and development of the .

The 3 Types of Testosterone Injections: Which is the Most . - GoodRx

Testosterone Injections: Types and Differences



Evidence Synthesis Available evidence, though limited, suggests that SC testosterone therapy in doses

similar to those given via IM route results in comparable pharmacokinetics and mean serum testosterone levels.

Testosterone Therapy: Review of Clinical Applications | AAFP

BEST PRACTICES IN ENDOCRINOLOGY: RECOMMENDATIONS FROM THE CHOOSING WISELY CAMPAIGN	
<i>Recommendation</i>	<i>Sponsoring organization</i>
Do not prescribe testosterone or testosterone products to men contemplating or attempting to initiate pregnancy.	American Society for Reproductive Medicine
Do not prescribe testosterone to men with erectile dysfunction who have normal testosterone levels.	American Urological Association
Do not prescribe testosterone therapy unless there is laboratory evidence of testosterone deficiency.	American Society for Clinical Pathology
Do not prescribe testosterone therapy unless there is biochemical evidence of testosterone deficiency.	The Endocrine Society/ American Association of Clinical Endocrinologists

Source: For more information on the Choosing Wisely Campaign, see <http://www.choosingwisely.org>. For supporting citations and to search Choosing Wisely recommendations relevant to primary care, see <http://www.aafp.org/afp/recommendations/search.htm>.

Depo-Testosterone has an average rating of 9.7 out of 10 from a total of 11 ratings on Drugs. 100% of reviewers reported a positive effect, while 0% reported a negative effect.

Depo-Testosterone, Aveed, Xyosted (testosterone cypionate, enanthate or .



Antares Pharmaceuticals' XYOSTED may be one of the best testosterone therapies available when evaluated against competitors such as Aveed, Depo-Testosterone, Delatestryl, and Testopel. Considering

- <https://telegra.ph/Buy-Deca-Durabolin-50-Mg-02-09>

- <https://telegra.ph/Genotropin-Pfizer-16-Iu-02-06>
- <https://telegra.ph/Pharma-Mix-4-Cycle-02-06>